arXiv:math/0111168v1 [math.AG] 14 Nov 2001

A strange example concerning Property N_p

Elena Rubei

Abstract

We exhibit an example of a line bundle M on a smooth complex projective variety Y s.t. M satisfies Property N_p for some p, the p-module of a minimal resolution of the ideal of the embedding of Y by M is nonzero and M^2 does not satisfy Property N_p.

Let M be a very ample line bundle on a smooth complex projective variety Y and let $\varphi_M : Y \to \mathbf{P}(H^0(Y,M)^*)$ be the map associated to M. We recall the definition of Property N_p of Green-Lazarsfeld (see [Gr1], [Gr2] and also [G-L], [Gr3]):
let Y be a smooth complex projective variety and let L be a very ample line bundle on Y defining an embedding $\varphi_L : Y \hookrightarrow \mathbf{P} = \mathbf{P}(H^0(Y,L)^*)$; set $S = S(L) = Sym^* H^0(L)$, the homogeneous coordinate ring of the projective space $\mathbf{P}$, and consider the graded S-module $G = G(L) = \oplus_n H^0(Y, L^n)$; let E_*

$$0 \longrightarrow E_n \longrightarrow E_{n-1} \longrightarrow \dots \longrightarrow E_0 \longrightarrow G \longrightarrow 0$$

be a minimal graded free resolution of G; let $E_p = \oplus_{q \geq 0} B_{p,q} \otimes S(-q)$; the line bundle L satisfies Property N_p ($p \in \mathbf{N}$) if and only if

$B_{0,0} = \mathbf{C}$
$B_{0,q} = 0$ *for* $q > 0$
$B_{i,q} = 0$ *for* $q \neq i+1$ *and* $1 \leq i \leq p$.

(Thus L satisfies Property N_0 if and only if $Y \subset \mathbf{P}(H^0(L)^*)$ is projectively normal, i.e. L is normally generated; L satisfies Property N_1 if and only if L satisfies Property N_0 and the homogeneous ideal I of $Y \subset \mathbf{P}(H^0(L)^*)$ is generated by quadrics; L satisfies Property N_2 if and only if L satisfies Property N_1 and the module of syzygies among quadratic generators $Q_i \in I$ is spanned by relations of the form $\sum L_i Q_i = 0$, where L_i are linear polynomials; and so on.)

Address: Elena Rubei, Dipartimento di Matematica “U. Dini”, via Morgagni 67/A 50134 Firenze, Italia

E-mail address: rubei@math.unifi.it

2000 Mathematical Subject Classification: 14C20, 13D02.

In [Ru] we considered the problem to see "how Property N_p propagates through powers": let M be a line bundle on a smooth complex projective variety Y and suppose M satisfies Property N_p; let $s \in \mathbf{N}$; we wondered for which k the line bundle M^s satisfies Property N_k. We proved: *Let Y be a smooth complex projective variety and M a line bundle on Y; if M satisfies Property N_p then M^s satisfies Property N_p if $s \geq p$.*
Besides, despite there are several theorems suggesting that there is a "propagation property" for Property N_p, we observed that surprisingly it is not true that if M satisfies Property N_p then any power of M satisfies Property N_p; the counterexample we exhibited was the following: the line bundle $M = \mathcal{O}_{\mathbf{P}^2}(2)$ satisfies Property N_p for every p (see the paper [O-P] and the references given there), but, by Ottaviani-Paoletti's Theorem in [O-P], for instance the bundle $M^2 = \mathcal{O}_{\mathbf{P}^2}(4)$ doesn't satisfy Property N_{10}.
The example can be considered not convincing by someone because the module E_{10} of a minimal resolution of $G(\mathcal{O}_{\mathbf{P}^2}(2))$ on $S(\mathcal{O}_{\mathbf{P}^2}(2))$ is 0.
The product of projective spaces furnishes the following more convincing example.

Example. Let $r \in \mathbf{N}$ and $r \geq 1$.
The line bundle $\mathcal{O}_{P^r \times P^1}(1,2)$ satisfies Property N_p $\forall p$ (see [O-P], Theorem 3.6).
On the contrary $\mathcal{O}_{P^r \times P^1}(2,4)$ does not satisfy Property N_{10}, in fact:
$\mathcal{O}_{P^1 \times P^1}(2,4)$ does not satisfy Property N_{10} by the following Gallego-Purnapranja's Theorem

`Theorem (Gallego-Purnapranja [G-P]).` *Let $a, b \geq 2$. The line bundle $\mathcal{O}_{\mathbf{P}^1 \times \mathbf{P}^1}(a,b)$ satisfies Property N_p if and only if $p \leq 2a + 2b - 3$.*

and the fact that $\mathcal{O}_{P^1 \times P^1}(2,4)$ does not satisfy Property N_{10} implies $\mathcal{O}_{P^r \times P^1}(2,4)$ does not satisfy Property N_{10}, by Remark in Section 2 of [Gr2] (precisely: write $\mathbf{P}^r = \mathbf{P}(V^*)$ and write the module E_p of a minimal resolution of $G(\mathcal{O}_{P^r \times P^1}(2,4))$ as $E_p = \oplus_{q \geq 0} B_{p,q} \otimes S(-q)$, as in the definition of Property N_p; we have that $B_{p,q}$ are representations of $GL(V)$; thus, if they are nonzero for $r = n$ then they are nozero also for $r > n$) (analogously to the proof of Prop. 1.8 in [O-P]).
If r is sufficiently great, we have that the 10-th module of a minimal resolution of $G(\mathcal{O}_{P^r \times P^1}(1,2))$ is nonzero (in fact, if X is a projective variety in $\mathbf{P}^n$ and $\dim X = m$ with $m < n - 1$, then the lenght l of a minimal resolution of the ideal $\mathcal{I}_X$ of X is $\geq n - m$: let t be s.t. $H^m(\mathcal{O}_X(t)) \neq 0$; thus $H^{m+1}(\mathcal{I}_X(t)) \neq 0$; then by breaking the twist by t of a minimal resolution of $\mathcal{I}_X$ into short exact sequences, we get $l \geq m - n$).

Acknowledgements. I thank G. Ottaviani for a helpful discussion.

References

[G-P] F.J. Gallego, B.P. Purnaprajna *Some results on rational surfaces and Fano varities* preprint math.AG/0001107

[Gr1] M. Green *Koszul cohomology and the geometry of projective varieties I* J. Differ. Geom. **20**, 125-171 (1984)

[Gr2] M. Green *Koszul cohomology and the geometry of projective varieties II* J. Differ. Geom. **20**, 279-289 (1984)

[Gr3] M. Green *Koszul cohomology and geometry*, in: M. Cornalba et al. (eds), Lectures on Riemann Surfaces, World Scientific Press (1989)

[G-L] M. Green, R. Lazarsfeld *On the projective normality of complete linear series on an algebraic curve* Invent. math. **83**, 73-90 (1986)

[O-P] G. Ottaviani, R. Paoletti *Syzygies of Veronese embeddings* preprint math.AG/9811131 to appear in Compositio Mathematica

[Ru] E. Rubei, *A note on Property* N_p, Manuscripta Mathematica 101 (2000), 449-455.